



Office of the State Long-Term Care

OMBUDSMAN

Established within the Iowa Department on Aging

Iowa Office of the State Long-Term Care Ombudsman

Volunteer Ombudsman Training Module

The Aging Process

2015

The mission of the Office of the State Long-Term Care Ombudsman is to protect the health, safety, welfare, and rights of individuals residing in long-term care by investigating complaints, seeking resolutions to problems, and providing advocacy with the goal of enhancing quality of life and care.

Table of Contents

Training Module: The Aging Process	3
SECTION 1: What is Aging?	4
SECTION 2: Biological Aspects of Aging	4
Structural	4
Sensory	5
Systems	8
SECTION 3: Psychological Aspects of Aging	9
SECTION 4: Sociological Aspects of Aging	11
SECTION 5: Myths and Stereotypes	14
SECTION 6: Common Illnesses and Conditions Associated with Aging	16
SECTION 7: Drugs and Their Side Effects in the Older Population	18
Module Quiz: The Aging Process	22
APPENDIX A: What is Aging?	23
APPENDIX B: A Profile of Older Americans: 2012	26
APPENDIX C: The Imperative for Good Care	44
APPENDIX D: Dementia	50
APPENDIX E: Alzheimer's Disease	52
APPENDIX F: Answer Key to Module Quiz	56

Instructions for Volunteer Ombudsmen

This training module is your "take home activity" to complete after attending Volunteer Ombudsman Training. The completion of this module is part of your certification requirements to become a Volunteer Ombudsman. Please review the materials in this module and complete the quiz, which can be found on page 22 of this module. The quiz can be completed while referring to the module materials. Record your score using the answer key found in Appendix F. After completing this module, you will have a chance to discuss your learnings with the Volunteer Ombudsman Coordinator during your post-training phone consultation.

Frequently used acronyms:

AoA – Administration on Aging

CMS – Centers for Medicare & Medicaid Services

LTC – Long-Term Care

LTCO – Long-Term Care Ombudsman

LTCOP – Long-Term Care Ombudsman Program

NCCNHR – National Citizens Coalition for Nursing Home Reform (now the National Consumer Voice for Quality Long-Term Care, or the “Consumer Voice”).

NORC – National Ombudsman Resource Center

OAA - Older Americans Act

Regs – Regulations

SLTCO – State Long-Term Care Ombudsman

SLTCOP – State Long-Term Care Ombudsman Program

Training Module: The Aging Process¹

Please feel free to complete the quiz questions as you explore this module. The quiz questions can be found on page 22.

Overview: This module provides basic information about the processes that occur throughout life, and particularly in the later years, that are considered normal aging. It also discusses common illnesses in later life and the effects of medications. Volunteer Ombudsmen must be able to work with older individuals and avoid stereotypes. Volunteer Ombudsmen should be alert to the difference between the effects of normal aging and the results of diseases that afflict some older persons.

Learning Objectives: At the conclusion of this module, Volunteer Ombudsmen will know:

- Normal age related changes;
- Myths and stereotypes about aging;
- Myths and stereotypes about care;
- Common illnesses and treatments; and
- The role of LTCO when poor care practices are encountered.

OPTIONAL: Before you start working your way through the module, take the brief interactive quiz linked below to determine how much you already know about the aging process. The quiz contains scenarios and links to fact sheets that you can refer to as you continue to read through this module.

What's Your Aging IQ?

(<http://www.niapublications.org/quiz/index.php>)

¹ Excerpted from The New Mexico Ombudsman Curriculum developed by Sara S. Hunt.

SECTION 1: What is Aging?

To better understand the population of long-term care residents who are a Volunteer Ombudsman's primary focus, you need to understand the "big picture" of the older population, defined by the OAA as age 60 or older. This section provides a profile of the older population in America today.

Please review the following documents:

- **What is Aging?** (See Appendix A)
 - **A Profile of Older Americans: 2012** (See Appendix B)
-

SECTION 2: Biological Aspects of Aging

Aging brings about some changes in all people. These changes are continuous throughout life, from losing baby teeth to the loss of taste buds. The normal changes with advanced age have only recently been studied and are beginning to be understood. Some changes are obvious in the way they alter physical appearance or in their visible effect upon body systems. Other changes are less apparent, in that they affect internal body systems, such as the circulatory systems. These changes vary in degree and rate from individual to individual.

Structural

MUSCLES

Muscles lose mass and tone. While exercise helps to maintain strength and tone, it does not prevent some loss. This change is observable in the looseness of underarm skin, sagging breasts, and thinner legs and arms reflecting the changes in musculature.

SKELETON

Another change affecting appearance is the flattening of the spongy "cushion" between the vertebrae. Over the years, this material loses its resiliency. Older people may be shorter than they were in younger years and have a stooped posture.

SKIN

There are several changes that affect the skin.

- The skin loses some elasticity, which results in wrinkles. The skin does not stretch and conform to its original shape as it once did.

- There is a loss in the natural oils in the skin, which may lead to dryness and scratchiness. Individuals may need to use moisturizer to replace the loss in oils.
- The skin becomes thinner and thus more susceptible to being broken or cut.
- Older people may become more sensitive to temperature changes.
- Some individuals may develop "aging" spots, which are dark areas of pigmentation. The presence of such spots does not indicate a problem with the function of the liver. The spots are simple changes in the pigmentation of the skin. Creams do not remove the spots although they may temporarily camouflage them. Spots on the skin of older people should be closely observed for sudden growth or changes in appearance. Such changes should be reported to a physician.

Sensory

MOUTH

The bone structure of the jaws may change, which can alter the way dentures fit. It is possible for an individual to develop problems with a set of dentures that he/she has had for years. Problems with dentures may have a negative impact on a person's nutritional intake.

- **TASTE:** The sensitivity of taste buds decreases with age, especially with men. The tastes that decline first are sweet and salty, with bitter and sour decreasing more slowly. Those changes mean that foods may not taste like they used to older people. Older individuals may over season food or may accuse others of omitting all seasonings in food preparation. Changes in taste may lead to a loss of appetite, which can lead to nutritional deficiencies.
- **SMELL:** Sensitivity to smell decreases as individuals age. Older individuals may be less aware of certain odors, even body odors, than younger people. The decreased sensitivity to smell may also adversely affect appetite.

VISION

There are several eye disorders that occur more often in the aged, such as glaucoma and cataracts. In the fourth decade of life, visual capacity begins to decline

- **DISTANCE:** The lens of the eye may lose some of its ability to accommodate changes in distance vision. That means that it may take a person a few seconds longer to recognize

someone who is across the room when the older person has been reading or doing handwork.

- **LIGHT:** The pupil of the eye tends to become smaller with age, permitting less light to enter the eye. This means eyes have a decreasing ability to adjust to changing amounts of light, and glare becomes a problem. Older people need more light than younger people do.² If an older person has been sitting in a semi-dark room and opens a door to find a visitor standing in bright sunlight, the older person may not immediately recognize the visitor. That does not indicate a problem with mental alertness, but it may indicate a longer than usual period of time required to adjust to differences in light.
- **COLOR:** Other changes in the lens of the eyes may make it difficult to distinguish blues and greens or pinks and yellows. An older person may comment on her green dress when it is actually blue. That kind of mistake does not necessarily indicate declining mental abilities; it may indicate changes in color identification. Colors that are very similar in shade like beige and brown may be difficult for older individuals to distinguish. Contrasting colors such as black and white may be more readily identified. Clothing can be tagged so those older individuals know which colors are complimentary.
- **DEPTH:** Changes in the eyes may affect an older person's mobility. The floor may appear to be rolling so that older people may shuffle along to ensure stable footing. Changes in depth perception can make it difficult to judge the height of curbs or steps. A person may take a large step and receive a jolt. It is helpful to edge steps or curbs in a bright, contrasting color to facilitate the older person's ability to distinguish distances and surface areas.
- **PRINT:** The lens of the eyes also lose some of its ability to focus on small print, such as the body of a newspaper. Headlines are more readily discernible. That means many of the forms that have instructions in small print are very difficult for older people to complete. The same is true of reading the statements of benefits, an activity schedule, a list of resident rights, or learning to operate the control knobs on a piece of equipment.

The cumulative effect of these vision changes can alter a person's sense of independence and self-confidence. If vision changes make it difficult for older individuals to negotiate a “strange” or unfamiliar environment, that person may limit shopping or take trips less often. An older person may appear to be two different people. One who is very efficient, steady, and independent may be observed in her own environment. In an unfamiliar environment, the same individual may appear

² Stuen, C., & Faye, E.E. “Vision Loss: Normal and Not Normal Changes among Older Adults.” *Generations*. XXVIII(1), p. 8.

confused, disoriented, and slow. That kind of difference may be due to vision changes. In the familiarity of a home environment, the person may function very well because he/she knows where everything is and how to operate the appliances.

It is important to allow older people the extra seconds needed for their eyes to accommodate to changes in light or distance. Eye examinations are also important to ensure that eye diseases or impairments are detected and promptly treated. Vision rehabilitation can be helpful in detecting problems and in offering tips to increase independent functioning.

HEARING

Changes in hearing are multiple and can have a profound effect upon the life of an older person. Hearing loss can cause depression and social isolation. Because it can lead to paranoia and suspicion, hearing loss is potentially the most problematic of perceptual losses. Individuals who have some degree of hearing loss may not realize that they have a loss.

When an individual with a hearing loss is in a group, the person with the hearing loss may begin to think that others are talking about him/her, or are deliberately excluding that person from the conversation. In reality, group members may not realize the need to face the person and to speak so that he/she follows the conversation. Individuals with hearing losses may hear part of what is said and not know they have heard only part of the statement or question.

The mind may automatically compensate for unintelligible conversation by inserting information, which seems to make sense. The person may then give an inappropriate response and not realize that the communication has been misunderstood.

There are three major types of hearing loss.

- **High frequency loss:** low, deep sounds are more readily heard than higher sounds.
- **Conductive hearing loss:** sound waves are not properly conducted to the inner ear making sounds become muffled and difficult to understand.
- **Central hearing loss:** allows speech to be heard but not understood. Signals from the ear either do not reach the brain or the brain misinterprets them.

Systems

CIRCULATORY SYSTEM

The heart, like other muscles, weakens and loses pumping capacity. Arteries or veins may become rigid or blocked, which restricts blood flow and circulation. Under routine circumstances, these changes do not greatly alter the daily functioning of an individual. These changes may be observed when an older person who has been sitting for a while suddenly stands and walks across the room. Unless a few extra seconds are allowed for the heart to supply sufficient blood to all the body extremities, the person may stumble, fall, or seem confused. After the heart has had sufficient time to pump the blood throughout the body, the unsteadiness or confusion disappears.

DIGESTIVE SYSTEM

One of the systems least affected by aging is the digestive system. As in earlier years, diet and exercise are extremely important to maintain proper functioning. Teeth become more brittle. Saliva, necessary to swallow food, decreases; the thirst response decreases. Peristalsis (the movement of the intestines) is slower, decreasing speed and effectiveness of digestion and elimination. Choking on food is a greater risk because of a decreased gag reflex.

URINARY SYSTEM

The urinary system experiences several changes.

- A general weakening of the bladder muscles means that the impulse to urinate cannot be delayed as long as in earlier years. When an older person says, "I have to go to the bathroom," that usually means now.
- The bladder doesn't stretch to hold as much as it used to, so urination may be more frequent. With weakened muscles the bladder may not empty completely which increases susceptibility to urinary infections.
- The kidneys filter the blood more slowly than in younger years. As a result, medications remain in the bloodstream longer than they do in younger people. The changes in functioning compounds the danger of over-medication. Dosages of medicine need to be closely and continuously monitored. Interaction effects between prescribed medicine and over-the-counter drugs, even aspirin or Bufferin, are more likely to occur.

REPRODUCTIVE SYSTEM

In the reproductive system there is little change. Vaginal secretions diminish; erections may require more stimulation. In men, the prostate may become enlarged. Regular check-ups are particularly important for men. Prostate trouble may go untreated until it requires radical treatment.

Summary

The cumulative effect of these changes is minimal in everyday functioning. These changes occur gradually, which allows individuals to adapt to the changes. Normal, daily functioning continues. The impact of these changes is more apparent when an older person is in an unfamiliar environment or when an older person is subjected to physical or psychological stress. Exercise and diet significantly impact the rate of these changes by slowing down the processes. In spite of the normal, age-related changes, older people function well enough to maintain daily functioning.

SECTION 3: Psychological Aspects of Aging

Memory

Short-term memory seems to decrease. It becomes more difficult to remember events in the immediate past, like what a person ate for breakfast, who came to visit yesterday, or the date and time of an appointment. There are ways to compensate for any decreases in short-term memory function. A person may write notes, which serve as reminders if they are kept in a specific place.

Freedom from distractions or too much stimulation may also help with remembering immediate events or information. Long-term memory seems to improve with increasing age. Events, which occurred forty or fifty years ago, may become easier to remember. As events are remembered and retold, they become more vivid and detailed.

Adaptation to Change

Everyone throughout their lives experiences change. When a person becomes older, he/she has lived through numerous changes. They have gone from the early days of automobiles to multi-lanes of traffic on interstates to airplanes to space ships. Individuals who have witnessed those changes have established patterns of adjusting to change. They know better what they can and can't tolerate and what is important to them.

Reactions to change vary from person to person. Change, whether positive or negative, is stressful. All individuals need time to adjust. Sometimes older people are seen as resistant to change, or "set in their ways." It may be that their refusal to accept change is a way of maintaining control. To say, "No," is to keep one area of their lives stable. At other times, change may be refused because it may not be understood. They may need more information or a clearer explanation, even if it is about a service being offered. Older people may need more time to consider the proposed change—to think it through, to decide. They may need assurance that the change can be tried on a temporary basis and then reevaluated. They may need reassurance about the terms of a service, information about other people who have utilized the service, and that the service can be easily

terminated, before he/she accepts the service. There may be a very good reason for saying, "No." They need to be listened to in order to understand their needs. Sometimes it is tough to find a balance between trusting their own priorities and understanding the enabling supports that they need.

Reminiscence

One method of coping with change is through reminiscence. There are several positive benefits of engaging in reminiscence. The present may be depressing or very unsatisfactory. By recalling a happier time, an older person may derive some contentment or the ability to endure the present. The strength to adjust to change may be derived from remembering previous successful adjustments.

Furthermore, reminiscence may provide an emotional outlet. Everyone reminisces. When something good happens; most people share the event with two or three friends. When friends meet, they sometimes recall previous shared experiences and relive them at that moment. Some older people may not have several different people with whom to share an experience. If only one or two people are around that older person, those individuals may hear the same story several times. Some of the common psychological purposes that reminiscence may serve are listed below.

- **IDENTITY:** Through storytelling, an older person can reveal personal achievements and characteristics. Indirectly, the older person may be saying, "*This is how I was before I became old.*" It serves as an introduction to that person prior to any limitations on energy or functioning. Personal characteristics are often revealed; a new acquaintance can begin to understand what the older person has been throughout his/her life by listening to reminiscences.
- **SELF ASSESSMENT:** In recalling the past, an older person may engage in self-assessment, deciding what kind of life one has lived. A review of the totality of one's life imparts a sense of integration of self. Allowing an older person to give advice, wisdom, or history to others through reminiscence can reinforce self-esteem. It may reinforce a person's feeling that his/her life has been worthwhile.
- **GRIEVING:** Reminiscence can be a productive method of dealing with loss and grief. In verbally sharing the loss, an individual may come to accept it. In grief, there is a need to remember and to relive past experiences. Reminiscence provides that opportunity. There may be conflicts in the past that are unresolved or need to be re-evaluated. By remembering past events, a person may decide to make amends with someone; to be forgiving or to seek forgiveness. Losses, which were suppressed, may surface. Grieving may need to be completed.

Intelligence

Intelligence does not decline with normal aging. When tested, older people scored lower on timed tests than do younger people. On tests without time limits, older people score better than younger individuals.

SECTION 4: Sociological Aspects of Aging

As with individuals of any age, familial relationships are important to older people. With increasing age, family composition often undergoes some changes. Older men are much more likely to be married than older women. Almost half of all older women are widows. Divorced and separated older persons represent only 10% of all older persons. However, this percentage has increased since 1980³.

Family connections extend into later life as reflected by living arrangements. Almost 60% of older women and 78% of older men live with a spouse or with another relative⁴.

Relationship patterns which were established in earlier years prevail into later life. If a parent and child have always had personality clashes, they will continue to unless they learn new ways of dealing with each other. The parent who listened primarily to one child or turned to a child for advice will continue that pattern unless something intervenes.

Role Reversal

While it is true that an older person may become more dependent in some capacities, the person is still an adult. Sometimes individuals may appear to act like children because they feel they are being treated as children especially when living in an institution. An individual may need transportation and assistance in completing forms. That does not mean that person needs someone to make financial decisions for him/her. An older individual may require temporary assistance in managing personal affairs until that person recovers from an illness or stress and is able to resume total responsibility. Sometimes families decide an older person is incapable of independence because the person makes a decision that disregards their advice.

Older people need to be encouraged to do as much for themselves as possible. Caregivers need to patiently allow sufficient time for persons to respond to questions or accomplish tasks. The emphasis should not be on perfection but on personal accomplishment. Volunteer Ombudsmen

³ *Profile of Older Americans 2003*, op.cit.

⁴ *Older Americans 2000: Key Indicators of Well-Being*. Appendix A: Detailed Tables. Indicator 5, Living Arrangements. Federal Interagency Forum on Aging-Related Statistics. <http://agingstats.gov>

should reinforce the decision-making ability of older individuals and expect and support as much independence in as many areas as possible.

Crisis

In families, it is helpful to anticipate potential crises. Before a stressful situation develops, consider the possibility that it may occur, and explore the alternatives. Areas to discuss include living arrangements, health care wishes, finances, wills, and funeral arrangements. It may be helpful to mention the subject and then discuss it more fully at a later date. Prior discussion helps prepare mental strategies for resolving crisis situations. It is easier to make decisions when everyone's wishes are known.

Limitations

Family relationships may involve some guilt. The guilt may be unjustified or due to unreasonable expectations. A personal re-assessment with realistic goals may be needed. If family members or an older relative makes excessive demands, a family conference or a one-on-one discussion may be in order. Problems, limitations, expectations, and responsibilities must be discussed. The older relative should be involved in the discussion and in problem solving. A workable solution must be found.

Losses

We experience losses throughout our lives. Some losses are more difficult to overcome than others. Common losses include the loss of friends, relatives, objects, and opportunities. Objects that are representative of special relationships or of personal achievement may be particularly important to an older person. Physical abilities may be lost: the use of an arm or leg, eyesight may diminish, and/or manual dexterity may decrease. These losses are usually accompanied by losses in roles and activities. The activities or functions which once gave meaning to one's life may have been dramatically altered. Opportunities to make new friends, acquire new skills, or accomplish lifelong goals, may be gone or greatly restricted. Recovery from losses may not be as quick in late life as it is in younger years.

There are two primary reactions to loss: anger and grief. Both are natural and may be expressed in various ways, depending on the individual. Talking about the loss is a therapeutic way to come to terms with it, to grieve, and accept the loss.

Death

Although death and dying may trigger strong feelings, it is a natural part of the life cycle. There are five major reactions to death or dying, which have been identified by researchers: denial, anger, bargaining, depression, and acceptance. Individuals do not always experience every stage, nor do they always experience the stages in the order listed. Stages may be repeated or skipped. Families or friends of a dying individual may also experience these reactions, and may do so at different times than the individual.

Visits With Residents Who Are Dying⁵

Responses of a Dying Person	Emotion	Role of LTC Ombudsman and Caregiver
When the awareness of a serious or fatal illness comes, persons react with shock and denial: " <i>No, not me! It can't be me!</i> " "This is not really happening. Someone has made a mistake."	DENIAL	Listening is very important. The dying person may not talk much and should not be pushed. Daydreaming about happier things, regardless of how improbable things may seem, should be encouraged and supported.
When denial can no longer be maintained, anger takes over. The question becomes " <i>Why me?</i> " or " <i>Why did God let this happen to me?</i> " The person feels angry, bitter, and envious of others who won't die.	ANGER	Family and friends usually find this stage difficult and mistake the anger as a personal attack. Be careful not to shorten or avoid visits or to react with anger. The resident needs an opportunity to ventilate his/her feelings. If the person feels respected and understood and is given attention by those important to him/her, she/he may soon begin to reduce the angry demands.
The person hopes that if she/he carries out promises, she/he will be rewarded with a longer life. This postponement is expressed in the hope that she/he will live to see some special event. " <i>Yes me, but...</i> " Many of these bargains are made with God and may be kept secret from family or friends.	BARGAINING	The resident needs someone to listen to him/her and to recognize his/her feelings. Expressing fears often helps to relieve the resident's feeling of guilt and enables the person to work through this stage in a more satisfying manner.

⁵ Adapted from Elisabeth Kubler-Ross

Responses of a Dying Person	Emotion	Role of LTC Ombudsman and Caregiver
Faced with the reality of such a great loss, the person is profoundly sad.	DEPRESSION	Our initial reaction to depression is to try to encourage the person to look at the bright side. This approach, however, can be an expression of our own needs and is not generally helpful in working through this stage. In dealing with reactive depression, the individual may have much to share. Listening is very important. The resident experiencing depression will often express his/her sorrow through silence. In these instances, a touch of a hand or just silently sitting together is usually more meaningful than words.
If the person has had sufficient time and the support and care of those around him/her, he/she will pass into a stage of acceptance of impending death: a calm, peaceful and comfortable readiness to face death. The person is not happy, but not terribly sad either.	ACCEPTANCE	The family may need more support than the dying person, who has already found some peace. It is a silent time in which the resident wishes to be left alone. He/she often prefers that visits be short and relatively silent. Our presence confirms that we will be around until the end and reassures him/her that he/she is not alone

SECTION 5: Myths and Stereotypes

Within American society, there are some common generalizations that are thought to be truths about older people. Many older people, who may expect these behaviors of themselves, believe these stereotypes. The myths, stereotypes, and negative attitudes greatly influence interactions with older people. Expectations about the later years are formed very early and are reinforced throughout life.

The truth is that there is great variety among individuals in later life. Individuals are what they have always been. There is as much diversity in personalities among older adults as there is among younger individuals. Problems arise when people act on their assumptions about the older person. Family members may unconsciously “watch” their older relatives to see when they will begin to exhibit these characteristics. Some major myths and stereotypes⁶ are listed below.

MYTH 1: Older people are disengaged.

Reality: Older people prefer to stay involved but their opportunities may be very limited.

MYTH 2: Older people are sick.

Reality: People do not become sick merely because they become older.

MYTH 3: Once a man, twice a child.

Reality: Older adults remain and function as adults.

MYTH 4: Older people are dependent.

Reality: Most older people are independent and take care of themselves.

MYTH 5: Older people are unproductive.

Reality: Most older people remain active and productive in their life.

MYTH 6: Older people become asexual.

Reality: Sexual desires continue throughout a person's life.

MYTH 7: Grandparents are always eager to be with their grandchildren.

Reality: Sometimes grandparents prefer visits that are planned in advance.

MYTH 8: Older people eventually become senile.

Reality: Older adults may experience cognitive decline, memory loss, or suffer from diseases such as Alzheimer's, but it is misleading to use catch all terms like "senile".

Volunteer Ombudsmen Need to Know!

Stereotyping and myths also affect the medical treatment older individuals receive and the way caregivers treat them. Clinical expertise is beginning to challenge many commonly held perceptions about inevitable age-related declines and appropriate interventions. As a Volunteer Ombudsman, you need to know which conditions indicate a need for more assessment and/or consideration of different treatment interventions instead of assuming that the conditions are simply manifestations of the aging process.

⁶ McFatter, J. 2012. http://prezi.com/cwuhakrkweg/?utm_campaign=share&utm_medium=copy&rc=ex0share

Since your role will be serving individuals in long term care facilities, this section will focus on applications in that environment. The same principles are applicable to individuals in home settings or other residences.

Please review the following document:

- **The Imperative for Good Care** (see Appendix C)
-

SECTION 6: Common Illnesses and Conditions Associated with Aging⁶

This section is included to provide basic information about selected conditions and illnesses that you might hear about as you visit residents. This information is not to be used as a medical guide.

Do not advise residents about treatment or make a diagnosis based on the following information.

Hiatus Hernia: Sixty-nine percent (69%) of people 70 years and older have hiatus hernias.

Hiatus hernias:

- Are protrusions of the stomach upward through the esophageal opening of the diaphragm.
- Can be somewhat minimized if the resident is sitting up straight while eating
- Are helped by smaller, more frequent meals as part of treatment
- Require the staff to realize the importance of positioning a person correctly

Constipation: The most common digestive problem among bedridden or inactive people is constipation.

Constipation can be caused by:

- Lack of fiber and fluid intake
- Decreased muscle tone
- Ignoring or being unable to heed the normal urge to defecate
- Laxative abuse
- Prolonged bed rest
- Insufficient food intake
- Tumors
- Certain medications, primarily tranquilizers, sedatives, pain medications, and antacids

Residents may complain about or have:

⁶ Excerpted and adapted from the Illinois Ombudsman Curriculum.

- Abdominal pain
- Distention of stomach
- Cramping

Many older people are dependent on laxatives. This dependency becomes counterproductive. If the person uses laxatives for any length of time, their digestive system will not function without them. Excessive use of laxatives impairs the absorption of fat and fat-soluble vitamins. Extreme constipation can become a medical emergency. It also can cause mental confusion as the system becomes poisoned by waste products that cannot be eliminated. However, a person who is dependent on laxatives needs to be taken off them slowly. A hearty breakfast, six or more glasses of liquid a day, and moderate exercise all are helpful in improving elimination.

Osteoporosis

Osteoporosis is:

- Loss of calcium from the bones
- Caused by insufficient calcium intake
- Lack of exercise
- Responsible for over 5 million spontaneous fractures every year; 55,000 people die annually from osteoporosis-related fractures. It is possible for bones to spontaneously break without being caused by a fall or applied pressure.
- Most prevalent in older white women

The vertebrae and other bones decrease in mass. This causes a gradual loss of height accompanied by a “dowager’s hump” (curving of the upper spine). Inactivity increases calcium depletion. Upon admission to a nursing facility, the older resident is generally less active than they would be in their home, which further accelerates the problem. Facility staff must include restorative nursing practices in resident daily routines including range of motion, standing and walking.

Dementia

Please review Appendix E.

Alzheimer's Disease

Please review Appendix F.

Parkinson's Disease

Parkinson’s is:

- A disease of the central nervous system
- Characterized by tremors in the extremities, rigidity, and slowness of movement
- An incurable, degenerative and progressive disease

Residents have:

- Tremor, which is a rhythmic shaking of a body part when it is at rest
- Poor grasp
- Poor mouth-hand coordination; the resident may need special utensils, special diets, and extended time to eat
- Rigidity or stiffness of muscles that may cause difficulty in walking, moving, or using one's arms and hands such as an inability to suck or close their lips well and limited ability to bite, chew and swallow
- Loss of balance and slowness of movement, as well as handwriting that gets smaller and smaller; loss of arm swing while walking
- Impassive facial expression
- Decreased volume and clarity of the person's voice

Tips for Volunteer Ombudsmen

Regardless of the cause of confusion, or whether it is reversible or irreversible, there are positive ways to respond to individuals. The expectation for improvement needs to be present. Individuals sometimes rise to meet our expectations, in spite of confusion. Voice tones as well as words and actions convey much meaning. As a Volunteer Ombudsman, you must be aware of all messages you are giving.

SECTION 7: Drugs and Their Side Effects in the Older Population⁷

Most long-term care residents are on five or more drugs at any time. Volunteer Ombudsmen, in visiting in nursing homes, will notice the side effects these drugs can have on residents. This section familiarizes Volunteer Ombudsmen with common drugs in long-term care facilities and the side effects many residents experience. Volunteer Ombudsmen should be familiar with this basic terminology of drugs so that when residents'/families' complaints involve drugs, Volunteer Ombudsmen recognize the terms and can thus refer the complaint reliably to the Local Ombudsman⁸.

Over a four year period, two-thirds of long-term care facility residents have adverse drug events (ADEs) and one out of seven of these results in hospitalization⁹.

⁷ Adapted from the Illinois Ombudsman Program Curriculum

⁸ For more information see Nguyen, C., and Williams, B. *Reducing Medication Problems in the Older*, USC School of Pharmacy 1995.

⁹ Cooper, JW. *Adverse Reaction Related Hospitalization of Nursing Facility Patients: A Four Year Study*, South Med Journal, 1999. 92:485-490.

The decision on prescribing appropriate drugs is the domain of the physician. Advance Practice Nurses (Nurse Practitioners and Clinical Nurse Specialists) as well as Physician Assistant may have prescriptive authority¹⁰. Pharmacists in long-term care facilities review the drug regime of residents on a monthly basis to ascertain if there are adverse drug reactions, allergies, contraindication, or ineffectiveness.

Remember: your role is not to second guess a medical decision regarding medications. You are to listen, observe, ask appropriate questions, and suggest that an individual ask his/her physician for additional review or more information. Volunteer Ombudsmen should know that all drugs given to an older person should be started at a low dose and raised slowly. This is especially true for individuals who have a dementia¹¹.

If more specific information related to medications is needed, call your Volunteer Coordinator or Local Ombudsman. You can also consult the following documents for excellent information about geriatric conditions, medications, and alternative treatments.

1. Neuroleptics (Major Tranquilizers, Anti-psychotics)

Used for psychosis, which is a severe mental disorder in which thinking and emotion are so impaired that the individual is seriously out of contact with reality. Examples of psychotic disorders include:

- **Schizophrenia**
- **Mania**
- **Depression with hallucinations**
- **Toxic psychosis** (LSD, PCP, etc.)
- **Organic brain syndrome** (infection, traumatic injury, tumors, degenerative disease, etc.)

Most common drugs used with these conditions:

- Haldol - Stelazine - Navene - Trilafon
- Loxitane - Moban - Prolixin - Thorazine

2. Minor Tranquilizers (Anti-Anxiety Agents)

Used for:

- Disabling anxiety (panic disorders, phobic disorders, post-traumatic stress disorder, and social phobia)
- Alcohol withdrawal
- Status epileptics
- Muscular spasms

¹⁰ *Ombudsman Guide to The Nursing Home Reform Law*, 2004, op. cit.

¹¹ *Resident Assessment Instrument for Long Term Care*, Centers for Medicare & Medicaid Services, Transmittal #8, Psychotropic /Drug RAP, 1999.

The drugs are grouped according to length of time it remains in a person's system:

- Long Life (18-36 Hr.), Valium, Paxipaim, Xanax (avoid these)
- Medium Life (10-12 Hr.), Tranxine, Ativan, Serax
- Short Life (4-12 Hr.), Librium, Centrax

Side effects:

- Common: Sedation
- Uncommon: Dry mouth, nausea, dizziness, confusion, withdrawal, tremor

Summary

All are potentially inappropriate for older persons, seldom habituating; usually used on short-term basis. Older persons are very sensitive to them.

3. Antidepressants

Used for:

Depression (when depression lasts more than two weeks):

- Norepinephrine type depression: Characterized by sleepiness, overeating, weight gain
- Serotonin type depression: Characterized by restlessness, anxiety, loss of appetite/weight
- Panic attacks
- Obsessive compulsive disorders: Where the mind is flooded with persistent and uncontrollable thoughts or is compelled to repeat an act again and again

Most common drugs used with these conditions:

- | | | | | |
|-----------|--------------|-----------|-------------|------------|
| - Asendin | - Imipramine | - Pamelor | - Sinequan | - Vivactil |
| - Desyrel | - Ludiomil | - Prozac | - Surmontil | - Zoloft |
| - Elavil | - Norpramin | - Paxil | - Tofranil | |

Common side effects:

- Sedation (more common with serotonergic)
- Dry mouth, nausea, constipation, sweats, and/or blurred vision

Summary

Very effective in endogenous depression; adverse reaction in people with cardiac problems and epilepsy; often requires two or three different medications before one is found that is effective and has the fewest side effects. IMPORTANT: These medications are toxic in overdoses. They can be quite dangerous. A small "mg" dosage is given for this reason.

4. Lithium Therapy

Used for:

- Bipolar disorders: Manic/depressive, manic, depression
- Alcoholism

Medication used:

- Lithium

Side effects:

- Common – Tremor, nausea, diarrhea
- Less common – Muscle weakness, muscle cramps, abdominal cramps, convulsions, acne, confusion

Summary

Most people take this medication without side effects. Therapy is long term (five years without a relapse). The patient is checked for blood levels and vital functions regularly.

5. Miscellaneous

1) Nasal decongestant sprays

Used for:

The relief of nasal congestion

Adverse Effects

- "Rebound congestion"
- Burning/stinging
- Sneezing

Examples:

- Dristan - Neo-Synephrine - Sinex

2) Caffeine

Used as an aid in staying awake

Found in several beverages like coffee, tea, colas, and cocoa

Adverse Effects of Caffeine

Insomnia, nausea/vomiting, restlessness, nervousness, excitement, increased urination, ringing in the ears

Quiz Questions

Training Module: The Aging Process

True or False

- T F 1. Depression in the older individuals is under diagnosed and under treated.
- T F 2. Medications prescribed for a medical problem may have an unintended effect on behavior.
- T F 3. It is possible for bones to break spontaneously.
- T F 4. Chronological age is an accurate indicator of an individual's feelings and abilities.
- T F 5. There are a range of non-drug interventions and treatments that may be effective in meeting a resident's needs when behavioral symptoms occur.
- T F 6. The possibility that a person will live in a care facility increases with age until age 78 when it remains the same no matter how long the person lives.
- T F 7. It is important to allow older people the extra seconds needed for their eyes to accommodate to changes in light or distance.
- T F 8. Individuals who have some degree of hearing loss may not realize that they have a loss.
- T F 9. One method of coping with change is through reminiscence.
- T F 10. Intelligence declines with the normal aging process.
- T F 11. "Once a man, twice a child," remains one of the great truths about the older.
- T F 12. Sensitivity to smells decreases with aging.
- T F 13. Sexual desire ceases in old age.
- T F 14. Movement is not important for individuals who are confined to bed.
- T F 15. Pressure sores are an unfortunate part of normal aging for frail, older persons.
- T F 16. Physical restraints prevent falls and injuries for individuals who are confused or have balance problems.
- T F 17. In spite of age related changes, individuals living in long-term care facilities are to be assisted in maintaining or improving their abilities unless a decline is unavoidable.
- T F 18. Inactivity increases calcium depletion which may contribute to osteoporosis.
- T F 19. Alzheimer's Disease affects areas of the brain that control long term memory first.
- T F 20. Incontinence is a normal part of aging.


APPENDIX A: What is Aging?

What is Aging

What is aging? Aging is a continuous process from birth to death, which encompasses physical, social, psychological, and spiritual changes.

Although aging is an ongoing process, the value of aging is seen differently at different points in the process. Some of the changes are anticipated with joy, such as a baby's first tooth or first step.

Other changes are greeted with a less positive response, such as pulling out the first gray hairs that appear. Youth is valued in American culture; while signs of aging are masked with face-lifts, wrinkle creams, and hair dyes. The process of physical maturation that is so eagerly anticipated in the first stages of life is viewed very negatively when the youthful attractiveness begins to change.



These prevailing attitudes lead to a denial of the signs of aging. Some individuals quit celebrating birthdays after a certain age. The stereotypical perceptions of aging as a period of deterioration and decline are therefore perpetuated. The positive aspects of aging are ignored. Each stage of life has its own pluses and minuses. Sometimes in old age, the balance may seem to tip to more negatives than positives, but this is not due to the *natural aging process*.

There are many positive aspects of aging. After 70 or 80 years of living, individuals tend to have a clear sense of their values and priorities. Older persons can make definite choices about how to use their time and energy. Their priorities may be very different from what caregivers, family, or friends want them to be. Older people have learned ways to adapt to changes; they have managed to survive. Advanced age can bring a freedom to speak one's opinion. Because of retirement, many older individuals have greater freedom to pursue interests, to use time to think and to reflect. To paraphrase Jung, as we age, we become more ourselves.


The advanced stages of aging are a normal, natural part of physical maturation. Instead of placing such a high value on youthfulness, it may be more productive to accept the changes throughout life without fear or denial.

B. Profile of Older People

As a long-term care ombudsman (LTCO), you will be working with older adults, their families, and their caregivers. To better understand the population of long term care residents who are your primary focus, you need to understand the "big picture" of the senior population, defined here as persons 65 years of age or older.

So who are aged people? At what age does a person become old? When a 64-year-old goes to bed and wakes up the next morning as a 65-year-old, has that person changed? Chronological age does not always correspond to a person's feelings. Although a person may be eighty years old, the person may feel like he/she is forty. The age a person feels may vary with the time of day, the day of the week, and/or activities or stresses present in that person's life. A person may

be very energetic on Saturday, but very tired and slow moving on Monday morning. Knowing a person’s chronological age tells you almost nothing about that individual’s feelings or abilities. Nevertheless, in this country, we categorize individuals by chronological age. Some key statistics¹ follow describing the population of seniors, persons 65 years or older.



Numbers and Growth

The older population—persons 65 years or older—numbered 35.6 million in 2002 (the most recent year for which data are available). They represented 12.3% of the U.S. population, about one in every eight Americans. The number of older Americans increased by 3.3 million or 10.2% since 1992, compared to an increase of 13.5% for the under-65 population. However, the number of Americans aged 45-64 – who will reach 65 over the next two decades – increased by 38% during this period.

The most rapid increase is expected between the years 2010 and 2030 when the “baby boom” generation reaches age 65. By 2030, there will be about 71.5 million older persons, more than twice their number in 2000. People 65+ represented 12.4% of the population in the year 2000 but are expected to be 20% of the population by 2030.

Minority Populations

Minority populations are projected to represent 26.4% of the elderly population in 2030, up from 17.2% in 2002. Between 2000 and 2030, the white** population 65+ is projected to increase by 77% compared with 223% for older minorities, including Hispanics, African-Americans,** American Indians, Eskimos, and Aleuts,** and Asians and Pacific Islanders.**

¹ The statistics and narrative information in this section come from: *A Profile of Older Americans 2003*, the Program Resources Department, American Association of Retired persons and the Administration on Aging, US Department of Health and Human Services, Washington, DC. <http://www.aoa.dhhs.gov/aoa/stats/profile/> The data is based on information from the US Bureau of the Census and the National Center for Health Statistics.


Age

The older population itself is getting older. In 2002, the 65-74 age group (18.3 million) was eight times larger than in 1900, but the 75-84 group (12.7 million) was more than 16 times larger and the 85+ group (4.6 million) was almost 38 times larger.

Living Arrangements

Over half of noninstitutionalized older persons lived with their spouse in 2002 (Figure 2). The proportion of individuals living with their spouse decreased with age, especially for women. **** About 30% of all older persons lived alone. The proportion living alone increases with advanced age. Among women aged 75 and over, for example, half lived alone (in 2000).

Figure 2: Living Arrangements of Persons 65+: 2002



Health and Health Care

In 2003, 38.6% of noninstitutionalized older persons assessed their health as excellent or very good, compared to 66.6% for persons aged 18-64. There was little difference between the sexes on this measure, but older African-Americans (57.7%) and older Hispanics (60.5%) were less likely to rate their health as excellent or good than were older Whites (75.4%).***** Most older persons have at least one chronic condition and many have multiple conditions. Among the most frequently occurring conditions of the elderly in 2000-2001 were: hypertension (49.2%), arthritic symptoms (36.1%), all types of heart disease (31.1%), any cancer (20.0), sinusitis (15.1%), and diabetes (15.0).

Nursing Homes

While a small number (1.56 million) and percentage (4.5%) of the 65+ population lived in nursing homes in 2000 the percentage increases dramatically with age, ranging from 1% for persons 65-74 years to 5% for persons 75-84 years and 18% for persons 85+.

A Profile of Older Americans: 2012



Administration on Aging
Administration for Community Living
U.S. Department of Health and Human Services

Table of Contents

Highlights	1
The Older Population	2
Future Growth	3
Figure 1: Number of Persons 65+, 1900 - 2060 (numbers in millions)	
Marital Status	4
Figure 2: Marital Status of Persons 65+ - 2012	
Living Arrangements	5
Figure 3: Living Arrangements of Persons 65+: 2012	
Racial and Ethnic Composition	6
Geographic Distribution	6
Figure 4: Persons 65+ as Percentage of Total Population by State: 2011 (US Map)	
Figure 5: Percentage Increase in Population 65+ by State: 2000 to 2011 (US Map)	
Figure 6: The 65+ Population by State: 2011 (table)	
Income	10
Figure 7: Percent Distribution by Income: 2011	
Poverty	11
Housing	12
Employment	12
Education	12
Health and Health Care	13
Health Insurance Coverage	14
Figure 8: Percentage of Persons 65+ with Health Insurance Coverage, 2011	
Disability and Activity Limitations	15
Figure 9: Percent of Persons with Limitations in Activities of Daily Living by Age Group: 2010	
Notes	16

Highlights*

- The older population (65+) numbered 41.4 million in 2011, an increase of 6.3 million or 18% since 2000.
- The number of Americans aged 45-64 – who will reach 65 over the next two decades – increased by 33% during this period.
- Over one in every eight, or 13.3%, of the population is an older American.
- Persons reaching age 65 have an average life expectancy of an additional 19.2 years (20.4 years for females and 17.8 years for males).
- Older women outnumber older men at 23.4 million older women to 17.9 million older men.
- In 2011, 21.0% of persons 65+ were members of racial or ethnic minority populations--9% were African-Americans (not Hispanic), 4% were Asian or Pacific Islander (not Hispanic), less than 1% were American Indian or Native Alaskan (not Hispanic), and 0.6% of persons 65+ identified themselves as being of two or more races. Persons of Hispanic origin (who may be of any race) represented 7% of the older population.
- Older men were much more likely to be married than older women--72% of men vs. 45% of women (Figure 2). 37% older women in 2012 were widows.
- About 28% (11.8 million) of noninstitutionalized older persons live alone (8.4 million women, 3.5 million men).
- Almost half of older women (46%) age 75+ live alone.
- In 2011, about 497,000 grandparents aged 65 or more had the primary responsibility for their grandchildren who lived with them.
- The population 65 and over has increased from 35 million in 2000 to 41.4 million in 2011 (an 18% increase) and is projected to increase to 79.7 million in 2040.
- The 85+ population is projected to increase from 5.7 million in 2011 to 14.1 million in 2040.
- Racial and ethnic minority populations have increased from 5.7 million in 2000 (16.3% of the elderly population) to 8.5 million in 2011 (21% of the elderly) and are projected to increase to 20.2 million in 2030 (28% of the elderly).
- The median income of older persons in 2011 was \$27,707 for males and \$15,362 for females. Median money income (after adjusting for inflation) of all households headed by older people rose by 2% (not statistically significant) from 2010 to 2011. Households containing families headed by persons 65+ reported a median income in 2011 of \$48,538.
- The major sources of income as reported by older persons in 2010 were Social Security (reported by 86% of older persons), income from assets (reported by 52%), private pensions (reported by 27%), government employee pensions (reported by 15%), and earnings (reported by 26%).
- Social Security constituted 90% or more of the income received by 36% of beneficiaries in 2010 (23% of married couples and 46% of non-married beneficiaries).
- Almost 3.6 million elderly persons (8.7%) were below the poverty level in 2011. This poverty rate is not statistically different from the poverty rate in 2010 (8.9%). During 2011, the U.S. Census Bureau also released a new Supplemental Poverty Measure (SPM) which takes into account regional variations in the livings costs, non-cash benefits received, and non-discretionary expenditures but does not replace the official poverty measure. The SPM shows a poverty level for older persons of 15.1% (more than 6 percentage points higher than the official rate of 8.7%). This increase is mainly due to including medical out-of-pocket expenses in the poverty calculations.

*Principal sources of data for the Profile are the U.S. Census Bureau, the National Center for Health Statistics, and the Bureau of Labor Statistics. The Profile incorporates the latest data available but not all items are updated on an annual basis.

The Older Population

The older population--persons 65 years or older—numbered 41.4 million in 2011 (the most recent year for which data are available). They represented 13.3% of the U.S. population, over one in every eight Americans. The number of older Americans increased by 6.3 million or 18% since 2000, compared to an increase of 9.4% for the under-65 population. However, the number of Americans aged 45-64 – who will reach 65 over the next two decades – increased by 33% during this period.

In 2011, there were 23.4 million older women and 17.9 million older men, or a sex ratio of 131 women for every 100 men. At age 85 and over, this ratio increases to 203 women for every 100 men.

Since 1900, the percentage of Americans 65+ has more than tripled (from 4.1% in 1900 to 13.3% in 2011), and the number has increased over thirteen times (from 3.1 million to 41.4 million). The older population itself is increasingly older. In 2011, the 65-74 age group (21.4 million) was almost 10 times larger than in 1900; the 75-84 group (12.8 million) was 16 times larger and the 85+ group (5 million) was 40 times larger.

In 2011, persons reaching age 65 had an average life expectancy of an additional 19.2 years (20.4 years for females and 17.8 years for males). A child born in 2011 could expect to live 78.7 years, about 30 years longer than a child born in 1900. Much of this increase occurred because of reduced death rates for children and young adults. However, the period of 1990-2007 also has seen reduced death rates for the population aged 65-84, especially for men – by 41.6% for men aged 65-74 and by 29.5% for men aged 75-84. Life expectancy at age 65 increased by only 2.5 years between 1900 and 1960, but has increased by 4.2 years from 1960 to 2007. Nonetheless, some research has raised concerns about future increases in life expectancy in the US compared to other high-income countries, primarily due to past smoking and current obesity levels, especially for women age 50 and over.

About 3 million persons celebrated their 65th birthday in 2011. In the same year, approximately 1.8 million persons 65 or older died. Census estimates showed an annual net increase between 2010 and 2011 of 916,837 in the number of persons 65 and over.

Between 1980 and 2010, the centenarian population experienced a larger percentage increase than did the total population. There were 53,364 persons aged 100 or more in 2010 (0.13% of the total 65+ population). This is a 66% increase from the 1980 figure of 32,194.


(Based on online data from the U.S. Census Bureau's 1) Population Estimates and Projections; 2) 2010 Census Special Reports, Centenarians: 2010, C2010SR-03, 2012; and 3) Table 5. Population by Age and Sex for the United States: 1900 to 2000, Part A. Hobbs, Frank and Nicole Stoops, Census 2000 Special Reports, Series CENSR-4, Demographic Trends in the 20th Century. The National Center for Health Statistics' Hoyert DL, Xu JQ. Deaths: Preliminary data for 2011. National vital statistics reports; vol 61 no 6. Hyattsville, MD: 2012. The National Research Council's Crimmins EM, Preston SH, Cohen B, editors. Explaining Divergent Levels of Longevity in High-Income Countries. Panel on Understanding Divergent Trends in Longevity in High-Income Countries, 2011.)

Future Growth

The older population will continue to grow significantly in the future (Figure 1). This growth slowed somewhat during the 1990's because of the relatively small number of babies born during the Great Depression of the 1930's. But the older population is beginning to burgeon as the "baby boom" generation begins to reach age 65.

The population 65 and over has increased from 35 million in 2000 to 41.4 million in 2011 (an 18% increase) and is projected to more than double to 92 million in 2060. By 2040, there will be about 79.7 million older persons, over twice their number in 2000. People 65+ represented 13.3% of the population in the year 2011 but are expected to grow to be 21% of the population by 2040. The 85+ population is projected to triple from 5.7 million in 2011 to 14.1 million in 2040.

Racial and ethnic minority populations have increased from 5.7 million in 2000 (16.3% of the elderly population) to 8.5 million in 2011 (21% of the elderly) and are projected to increase to 20.2 million in 2030 (28% of the elderly). Between 2012 and 2030, the white (not Hispanic) population 65+ is projected to increase by 54% compared with 125% for older racial and ethnic minority populations, including Hispanics (155%), African-Americans (not Hispanic) (104%), American Indian and Native Alaskans (not Hispanic) (116%), and Asians (not Hispanic) (119%).




Note: Increments in years are uneven.

(Based on online data from the U.S. Census Bureau's 1) Population Estimates and Projections; 2) Table 1. Projected Population by Single Year of Age (0-99, 100+), Sex, Race, and Hispanic Origin for the United States: July 1, 2012 to July 1, 2060, Release Date: 2012; and 3) Table 5. Population by Age and Sex for the United States: 1900 to 2000, Part A. Hobbs, Frank and Nicole Stoops, Census 2000 Special Reports, Series CENSR-4, Demographic Trends in the 20th Century.)

Marital Status

In 2012, older men were much more likely to be married than older women--72% of men, 45% of women (Figure 2). Widows accounted for 37% of all older women in 2012. There were over four times as many widows (8.5 million) as widowers (2.1 million).

Divorced and separated (including married/spouse absent) older persons represented only 12% of all older persons in 2012. However, this percentage has increased since 1980, when approximately 5.3% of the older population were divorced or separated/spouse absent.



(Based on online data from the U.S. Census Bureau's Current Population Survey, Annual Social and Economic Supplement.)

Living Arrangements


Over half (57%) the older noninstitutionalized persons lived with their spouse in 2012. Approximately 13.2 million or 72% of older men, and 10.3 million or 45% of older women, lived with their spouse (Figure 3). The proportion living with their spouse decreased with age, especially for women. Only 32% of women 75+ years old lived with a spouse.

About 28% (11.8 million) of all noninstitutionalized older persons in 2012 lived alone (8.4 million women, 3.5 million men). They represented 36% of older women and 19% of older men. The proportion living alone increases with advanced age. Among women aged 75 and over, for example, almost half (46%) lived alone.

In 2011, a total of about 2 million older people lived in a household with a grandchild present. About 497,000 of these grandparents over 65 years old were the persons with primary responsibility for their grandchildren who lived with them.

A relatively small number (1.5 million) and percentage (3.6%) of the 65+ population in 2011 lived in institutional settings such as nursing homes (1.3 million). However, the percentage increases dramatically with age, ranging (in 2011) from 1% for persons 65-74 years to 3% for persons 75-84 years and 11% for persons 85+. In addition, in 2009, approximately 2.7% of the elderly lived in senior housing with at least one supportive service available to their residents.

Figure 3: Living Arrangements of Persons 65+, 2012



(Based on online data from the U.S. Census Bureau's American Community Survey. The Centers for Medicare and Medicaid Services' Medicare Current Beneficiary Survey.)

Racial and Ethnic Composition

In 2011, 21.0% of persons 65+ were members of racial or ethnic minority populations--9% were African-Americans (not Hispanic), 4% were Asian or Pacific Islander (not Hispanic), less than 1% were American Indian or Native Alaskan (not Hispanic), and 0.6% of persons 65+ identified themselves as being of two or more races. Persons of Hispanic origin (who may be of any race) represented 7% of the older population.

Only 7.4% of all the people who were members of racial and ethnic minority populations were 65+ in 2011 (9.2% of African-Americans (not Hispanic), 5.7% of Hispanics, 9.8% of Asians and Pacific Islanders (not Hispanic), 8.4% of American Indians and Native Alaskans (not Hispanic)) compared with 16.7% of non-Hispanic whites.

(Based on online data from the U.S. Census Bureau's Population Estimates and Projections.)

Geographic Distribution


The proportion of older persons in the population varies considerably by state with some states experiencing much greater growth in their older populations (Figures 4 and 5). In 2011, over half (51%) of persons 65+ lived in 9 states: California (4.4 million); Florida (3.4 million); New York (2.7 million); Texas (2.7 million); Pennsylvania (2.0 million); and Ohio, Illinois, Michigan, North Carolina and each had well over 1 million (Figure 6).

Persons 65+ constituted approximately 15% or more of the total population in 11 states in 2011: Florida (17.6%); Maine (16.3%); West Virginia (16.2%); Pennsylvania (15.6%); Montana (15.2%); Arkansas (15%); Delaware (15%); Hawaii (15%); Iowa (15%); Rhode Island (15%); and Vermont (15.0%). In 13 states, the 65+ population increased by 30% or more between 2000 and 2011: Alaska (58%), Nevada (53%), Arizona (37%), Colorado (37%), Georgia (37%), Idaho (37%), South Carolina (35%), Utah (35%), New Mexico (33%), North Carolina (32%), Delaware (31%), Texas (30%), and Washington (30%). The 17 jurisdictions with poverty rates at or over 10% for elderly during 2011 were: Mississippi (13.5%), Louisiana (12.7%), District of Columbia (12.5%), Kentucky (11.8%), New Mexico (11.8%), New York (11.7%), Texas (11.4%), South Dakota (11.1%), Georgia (10.9%), Tennessee (10.7%), Arkansas (10.5%), North Dakota (10.5%), Alabama (10.3%), Rhode Island (10.2%), South Carolina (10.1%), California (10.0%), and Florida (10.0%) (Figure 6):

Most persons 65+ lived in metropolitan areas in 2011 (81%). About 66% of these older persons lived outside principal cities and 34% lived inside principal cities. Also, 19% of older persons lived outside of metropolitan areas.


The elderly are less likely to change residence than other age groups. From 2011 to 2012, only 3% of older persons moved as opposed to 14% of the under 65 population. Most older movers (61%) stayed in the same county and 83% remained in the same state. Only 16% of the movers moved from out-of-state or abroad.

Figure 4: Persons 65+ as a Percentage of Total Population, 2011



(Source: 2011 Population Estimates from the U.S. Census Bureau)

Figure 5: Percent Increase in Population 65+, 2000 to 2011



(Source: 2000 and 2011 Population Estimates from the U.S. Bureau of the Census)

Figure 6: The 65+ Population by State 2011


State	Number of Persons 65 and Older	Percent of All Ages	Percent Increase from 2000 to 2011	Percent Below Poverty 2011
US Total (50 States + DC)	41,394,141	13.30%	18.03%	8.7%
Alabama	672,586	14.00%	15.86%	10.3%
Alaska	58,213	8.10%	58.11%	6.3%
Arizona	921,835	14.20%	37.30%	8.5%
Arkansas	429,099	14.60%	14.56%	10.5%
California	4,398,624	11.70%	21.88%	10.0%
Colorado	575,820	11.30%	37.23%	7.8%
Connecticut	515,459	14.40%	9.47%	6.8%
Delaware	133,464	14.70%	30.73%	5.7%
District of Columbia	70,179	11.40%	0.65%	12.5%
Florida	3,360,195	17.60%	19.51%	10.0%
Georgia	1,077,512	11.00%	36.51%	10.9%
Hawaii	202,544	14.70%	25.23%	8.2%
Idaho	202,433	12.80%	37.28%	6.8%
Illinois	1,640,765	12.70%	9.23%	8.2%
Indiana	858,087	13.20%	13.64%	7.2%
Iowa	457,738	14.90%	4.82%	6.9%
Kansas	382,661	13.30%	7.26%	7.2%
Kentucky	591,851	13.50%	17.24%	11.8%
Louisiana	572,440	12.50%	10.87%	12.7%
Maine	216,218	16.30%	17.50%	8.0%
Maryland	729,488	12.50%	21.32%	7.5%
Massachusetts	921,889	14.00%	7.08%	9.3%
Michigan	1,389,155	14.10%	13.59%	8.2%
Minnesota	701,768	13.10%	17.56%	8.0%
Mississippi	388,071	13.00%	13.15%	13.5%
Missouri	854,652	14.20%	13.07%	8.2%
Montana	151,452	15.20%	24.97%	8.1%
Nebraska	250,382	13.60%	7.71%	7.8%
Nevada	339,478	12.50%	53.10%	9.4%
New Hampshire	184,376	14.00%	24.19%	6.2%
New Jersey	1,208,360	13.70%	8.48%	7.8%
New Mexico	282,375	13.60%	32.61%	11.8%
New York	2,664,694	13.70%	8.62%	11.7%
North Carolina	1,278,786	13.20%	31.67%	9.7%
North Dakota	98,595	14.40%	4.45%	10.5%
Ohio	1,648,444	14.30%	9.26%	7.7%
Oklahoma	517,654	13.70%	13.70%	9.5%
Oregon	553,061	14.30%	25.87%	7.4%
Pennsylvania	1,981,565	15.60%	3.31%	8.0%
Rhode Island	154,143	14.70%	1.16%	10.2%
South Carolina	658,561	14.10%	35.34%	10.1%
South Dakota	118,653	14.40%	9.81%	11.1%
Tennessee	879,651	13.70%	24.86%	10.7%
Texas	2,706,029	10.50%	30.21%	11.4%
Utah	259,221	9.20%	35.28%	5.3%
Vermont	94,122	15.00%	20.75%	7.0%
Virginia	1,011,063	12.50%	27.36%	7.5%
Washington	864,873	12.70%	30.15%	8.5%
West Virginia	300,970	16.20%	8.80%	9.4%
Wisconsin	792,620	13.90%	12.49%	7.5%
Wyoming	72,267	12.70%	24.33%	6.2%
Puerto Rico	600,927	15.00%	40.00%	39.00%

(Based on online data from the U.S. Census Bureau's 1) Population Estimates and Projections; 2) American Community Survey; 3) Current Population Survey, Annual Social and Economic Supplement; and 4) International Data Base.)


Income

The median income of older persons in 2011 was \$27,707 for males and \$15,362 for females. From 2010 to 2011, median money income (after adjusting for inflation) of all households headed by older people rose 2% but this was not statistically significant. Households containing families headed by persons 65+ reported a median income in 2011 of \$48,538 (\$50,658 for non-Hispanic Whites, \$33,809 for Hispanics, \$39,533 for African-Americans, and \$50,971 for Asians). About 5% of family households with an elderly householder had incomes less than \$15,000 and 67% had incomes of \$35,000 or more (Figure 7).

Figure 7: Percent Distribution by Income: 2011



\$48,538 median for 14.4 million family households 65+



\$19,939 median for 40.2 million persons 65+ reporting income.

For all older persons reporting income in 2011 (40.2 million), 17.8% reported less than \$10,000 and 40% reported \$25,000 or more. The median income reported was \$19,939.

The major sources of income as reported by older persons in 2010 were Social Security (reported by 86% of older persons), income from assets (reported by 52%), private pensions (reported by 27%), government employee pensions (reported by 15%), and earnings (reported by 26%). In 2010, Social Security benefits accounted for 37% of the aggregate income of the older population. The bulk of the remainder consisted of earnings (30%), asset income (11%), and pensions (18%). Social Security constituted 90% or more of the income received by 36% of beneficiaries (23% of married couples and 46% of non-married beneficiaries).

(Based on online data from the U.S. Census Bureau's 1) Current Population Survey, Annual Social and Economic Supplement; and 2) "Income, Poverty, and Health Insurance Coverage in the United States: 2011," P60-243, issued September, 2012. The Social Security Administration's "Fast Facts and Figures About Social Security, 2012.")

Poverty

Over 3.6 million elderly persons (8.7%) were below the poverty level in 2011. This poverty rate is not statistically different from the poverty rate in 2010 (8.9%). Another 2.4 million or 5.8% of the elderly were classified as "near-poor" (income between the poverty level and 125% of this level).

Just over 1.6 million older Whites (not Hispanic) (6.7%) were poor in 2011, compared to 17.3% of elderly African-Americans, 11.7% of Asians, and 18.7% of elderly Hispanics. Higher than average poverty rates were found in 2011 for older persons who lived inside principal cities (11.7%) and in the South (10.1%).

Older women had a higher poverty rate (10.7%) than older men (6.2%) in 2011. Older persons living alone were much more likely to be poor (16.5%) than were older persons living with families (5%). The highest poverty rates were experienced among older Hispanic women (38.8%) who lived alone and also by older Black women (32.2%) who lived alone.

In 2011, the U.S. Census Bureau released a new Supplemental Poverty Measure (SPM). The SPM methodology shows a significantly higher number of older persons below poverty than is shown by the official poverty measure. For persons 65 and older this poverty measure shows a poverty level of 15.1% in 2011 (more than 6 percentage points higher than the official rate of 8.7%). Unlike the official poverty rate, the SPM takes into account regional variations in the cost of housing etc. and, even more significantly, the impact of both non-cash benefits received (e.g., SNAP/food stamps, low income tax credits, WIC, etc.) and non-discretionary expenditures including medical out-of-pocket (MOOP) expenses. For persons 65 and over, MOOP was the major source of the significant differences between these measures. Bear in mind that the SPM does not replace the official poverty measure.

(Based on online data from the U.S. Census Bureau's 1) Current Population Survey, Annual Social and Economic Supplement; 2) "Income, Poverty, and Health Insurance Coverage in the United States: 2011," P60-243, issued September, 2012; and 3) "The Research Supplemental Poverty Measure," P60-244, issued November 2012.)

Housing

Of the 25.1 million households headed by older persons in 2011, 81% were owners and 19% were renters. The median family income of older homeowners was \$32,900. The median family income of older renters was \$16,200. In 2011, almost 50% of older householders spent more than one-fourth of their income on housing costs - 43% for owners and 71% for renters - as compared to 50% of all householders.

For older homeowners in 2011, the median construction year was 1970 compared with 1976 for all homeowners. Among the homes owned by people age 65 and older, 3.3% had physical problems. In 2011, the median value of homes owned by older persons was \$150,000 (with a median purchase price of \$55,000) compared to a median home value of \$160,000 for all homeowners. About 65% of older homeowners in 2011 owned their homes free and clear.

(Based on online data sponsored by the Department of Housing and Urban Development, available from the U.S. Census Bureau, American Housing Survey, National Tables: 2011.)

Employment

In 2012, 7.7 million (18.5 %) Americans age 65 and over were in the labor force (working or actively seeking work), including 4.3 million men (23.6%) and 3.4 million women (14.4%). They constituted 5% of the U.S. labor force. About 6.2% were unemployed. Labor force participation of men 65+ decreased steadily from 2 of 3 in 1900 to 15.8% in 1985; then stayed at 16%-18% until 2002; and has been increasing since then to over 20%. The participation rate for women 65+ rose slightly from 1 of 12 in 1900 to 10.8% in 1956, fell to 7.3% in 1985, was around 7%-9% from 1986 – 2002. However, beginning in 2000, labor force participation of older women has been gradually rising to the 2012 level. This increase is especially noticeable among the population aged 65-69.

(Based on online data from the Bureau of Labor Statistics' Current Population Survey, Labor Force Statistics.)

Education

The educational level of the older population is increasing. Between 1970 and 2012, the percentage of older persons who had completed high school rose from 28% to 81%. About 24% in 2012 had a bachelor's degree or higher. The percentage who had completed high school varied considerably by race and ethnic origin in 2012: 86% of Whites (not Hispanic), 74% of Asians, 69% of African-Americans, 69% of American Indian/Alaska Natives, and 49.0% of Hispanics. The increase in educational levels is also evident within these groups. In 1970, only 30% of older Whites and 9% of older African-Americans were high school graduates.

(Based on online data from the U.S. Census Bureau's Current Population Survey, Annual Social and Economic Supplement.)

Health and Health Care

In 2012, 44% of noninstitutionalized older persons assessed their health as excellent or very good (compared to 64% for persons aged 18-64 years). There was little difference between the sexes on this measure, but older African-Americans (not Hispanic) (25.8%), older American Indians/Alaska Natives (29%), older Asians (33%), and older Hispanics (29.7%) were less likely to rate their health as excellent or very good than were older Whites (not Hispanic) (44.7%). Most older persons have at least one chronic condition and many have multiple conditions. In 2009-2011, the most frequently occurring conditions among older persons were: diagnosed arthritis (51%), all types of heart disease (31%), any cancer (24%), diagnosed diabetes (20% in 2007-2010), and hypertension (high blood pressure or taking antihypertensive medication) (72 percent in 2007-2010).

In 2012, 68% of people age 65 and over reported that they received an influenza vaccination during the past 12 months and 60% reported that they had ever received a pneumococcal vaccination. About 28% (of persons 60+) reported height/weight combinations that placed them among the obese. Almost 45% of persons aged 65-74 and 29% of persons 75+ reported that they engaged in regular leisure-time physical activity. Only 9% reported that they are current smokers and 6% reported excessive alcohol consumption. Only 2% reported that they had experienced psychological distress during the past 30 days.

In 2010, about 13.6 million persons aged 65 and older were discharged from short stay hospitals. This is a rate of 3,326.4 for every 10,000 persons aged 65+ which is about three times the comparable rate for persons of all ages (which was 1,102 per 10,000). The average length of stay for persons aged 65-74 was 5.4 days; for ages 75-84 it was 5.7 days; and for ages 85 and over it was 5.6 days. The comparable rate for persons of all ages was 4.8 days. The average length of stay for older people has decreased by 5 days since 1980. Older persons averaged more office visits with doctors in 2011. Among people age 75 and over, 21 percent had 10 or more visits to a doctor or other health care professional in the past 12 months compared to 14 percent among people age 45 to 64. In 2012, almost 96% of older persons reported that they did have a usual place to go for medical care and only 2.4% said that they failed to obtain needed medical care during the previous 12 months due to financial barriers.


In 2011 older consumers averaged out-of-pocket health care expenditures of \$4,769, an increase of 46% since 2000. In contrast, the total population spent considerably less, averaging \$3,313 in out-of-pocket costs. Older Americans spent 12.2% of their total expenditures on health, almost twice the proportion spent by all consumers (6.7%). Health costs incurred on average by older consumers in 2011 consisted of \$3,076 (64%) for insurance, \$786 (16%) for medical services, \$714 (15%) for drugs, and \$193 (4.0%) for medical supplies.

(Based on online data from the National Center for Health Statistics' 1) Health Data Interactive data warehouse; and 2) Early Release of Selected Estimates Based on Data From the January–September 2012 National Health Interview Survey. The Bureau of Labor Statistics' Consumer Expenditure Survey.)

Health Insurance Coverage

In 2011, almost all (93%) non-institutionalized persons 65+ were covered by Medicare. Medicare covers mostly acute care services and requires beneficiaries to pay part of the cost, leaving about half of health spending to be covered by other sources. About 58% had some type of private health insurance. Over 9% had military-based health insurance and 9% of the non-institutionalized elderly were covered by Medicaid (Figure 8). Less than 2% did not have coverage of some kind. About 86% of non-institutionalized Medicare beneficiaries in 2009 had some type of supplementary coverage. Among Medicare beneficiaries residing in nursing homes, almost half (49%) were covered by Medicaid.

Figure 8:



Note: Figure 8 data are for the non-institutionalized elderly. A person can be represented in more than one category.


(Based on online data from the Centers for Medicare and Medicaid Services' Medicare Current Beneficiary Survey. The U.S. Census Bureau's "Income, Poverty, and Health Insurance Coverage in the United States: 2011," P60-243, issued September, 2012.)

Disability and Activity Limitations

Some type of disability (i.e., difficulty in hearing, vision, cognition, ambulation, self-care, or independent living) was reported by 35% of men and 38% of women age 65+ in 2011. Some of these disabilities may be relatively minor but others cause people to require assistance to meet important personal needs. There is a strong relationship between disability status and reported health status. Presence of a severe disability is also associated with lower income levels and educational attainment.

Using limitations in activities of daily living (ADLs) and instrumental activities of daily living (IADLs) to measure disability, 28% of community-resident Medicare beneficiaries age 65+ reported difficulty in performing one or more ADL and an additional 12% reported difficulty with one or more IADL. By contrast, 92% of institutionalized Medicare beneficiaries had difficulties with one or more ADLs and 76% of them had difficulty with three or more ADLs. [ADLs include bathing, dressing, eating, and getting around the house. IADLs include preparing meals, shopping, managing money, using the telephone, doing housework, and taking medication.] Limitations in activities because of chronic conditions increase with age. As shown in Figure 9, the rate of limitations in activities among noninstitutionalized persons 85 and older are much higher than those for persons 65-74.

Figure 9:



Except where noted, the figures above are taken from surveys of the noninstitutionalized elderly. Although nursing homes are being increasingly used for short-stay post-acute care, about 1.3 million elderly are in nursing homes (more than half are age 85 and over). These individuals often need care with their ADLs and/or have severe cognitive impairment due to Alzheimer's disease or other dementias.

(Based on online data from the U.S. Census Bureau's American Community Survey. The Centers for Medicare and Medicaid Services' Medicare Current Beneficiary Survey. The National Center for Health Statistics, including the NCHS Health Data Interactive data warehouse.)

Notes:

*Principal sources of data for the Profile are the U.S. Census Bureau, the National Center for Health Statistics, and the Bureau of Labor Statistics. The Profile incorporates the latest data available but not all items are updated on an annual basis.

A Profile of Older Americans: 2012 was developed by the Administration on Aging (AoA), Administration for Community Living, U.S. Department of Health and Human Services.

AoA serves as an advocate for the elderly within the federal government and is working to encourage and coordinate a responsive system of family and community based services throughout the nation. AoA helps states develop comprehensive service systems which are administered by 56 State and Territorial Units on Aging, 629 Area Agencies on Aging, 246 Native American and Hawaiian organizations, and approximately 20,000 local service providers.

APPENDIX C: The Imperative for Good Care

The Imperative for Good Care

In addition to challenging some of the long-held perceptions about the causes of decline and appropriate treatment, there is a solid legal basis for rethinking stereotypical responses.¹ The Nursing Home Reform Law (OBRA '87) challenges the mindset that “*this is the way we’ve always done it,*” or “*we don’t have the staff to do it.*”

OBRA challenges everyone to re-examine assumptions and practices: *that old people are hopelessly depressed; bedsores and incontinence are unavoidable, and residents must be restraints help residents.* There are practitioners² who have blazed the trail: finding that time spent on thorough assessment and care planning saves time in the long run; accommodating individual needs is possible and is more efficient; and eliminating restraints results in better care. Their experience shows the law’s potential.

One of the principle provisions of OBRA, Quality of Care, says, “*A nursing facility must provide services and activities to attain or maintain the highest practicable physical, mental, and psychosocial well-being of each resident in accordance with a written plan of care.*”

The requirements for long term care facilities explains what Quality of Care means:

“Based on a comprehensive assessment of a resident, the facility must ensure that a resident’s abilities in activities of daily living do not diminish unless circumstances of the individual’s clinical condition demonstrate that diminution was unavoidable. This includes the resident’s ability to bathe, dress, and groom; transfer and ambulate; toilet; eat; and use speech language or other functional communication systems.”

The regulation applies to vision and hearing, pressure sores, urinary incontinence, range of motion, mental and psychosocial functioning, naso-gastric tubes, and other areas of care.

In short, this provision means that people should not get worse because of what the nursing home does to them. In fact, they should reach the highest level of functioning and well-being that they are capable of achieving. If a resident was able to walk, transfer, bathe himself/herself, move his/her arms, or maintain his/her skin condition when he/she entered the facility, he/she should still be able to do so after six months or a year; actually, for the rest of his/her stay in the facility, unless circumstances of his/her clinical condition demonstrate that diminution or decline was unavoidable. There are only three reasons that diminution is unavoidable:

- A new disease or condition is experienced by a resident (e.g. heart disease added to the Parkinson’s)
- A resident’s disease progresses (e.g. the parkinsonian medicine no longer works and the individual becomes so rigid he is rendered immobile)
- A resident refuses care.

¹ Most of this section is from Frank, B., “The Promise of Nursing Home Reform Is In Your Hands . . . An Advocate’s Message to Surveyors,” *Survey and Certification Review*, June 1992. pp3-8, and the *Ombudsman Guide to The Nursing Home Reform Law*. National Long-Term Care Ombudsman Resource Center, National Citizens’ Coalition for Nursing Home Reform. Washington, DC. 2004.

² The Pioneer Network serves as a national focal point for this type of activity, promoting “culture change,” cultivating and sharing best practices. P.O. Box 18648, Rochester, NY 14618. (515)271-7570. www.pioneernetwork.net

The following is a description of some common myths and stereotypes that are being proved untrue. The Resident Assessment Protocols, part of the mandatory resident assessment process, contain excellent guidance to assist in changing perceptions and treatment approaches for all the conditions in this section. The knowledge basis and educational resources are available to alter *the way we've always done things*. As we change our way of thinking about conditions, there will be dramatic differences in what happens to individuals who enter nursing facilities.

LOSS OF MOBILITY

MYTH OR STEREOTYPE

Given the frail condition of residents, movement is not as important for them as it is for other adults. They will experience a decline in mobility as an inevitable part of growing older.

REALITY

“Movement, like other basic human needs, is lifelong and doesn’t end with [old age and] institutionalization. The ability to meet these needs may fluctuate with physical and mental ability, but the drive that initiates the pursuit is forever. Frail, elderly persons who enter nursing facilities retain the drive to meet their need for movement, just as they do for the other basic needs. Institutions often fail to assist residents in meeting movement needs because they fail to recognize movement as a basic human need.”³

All individuals need to move. “Impaired mobility can lead to a number of harmful physical and mental complications, which taken to their extreme, can be fatal.”⁴ Immobility negatively affects every body system. The effect of immobility, as well as ways to *maintain* mobility, is documented.

In a limited study of nursing home residents, those who walked outdoors reported less fatigue than residents who did not.⁵ Residents in the walking group slept better and reported better appetites than others in the study. Mobility is essential to life. It affects more aspects of life than just the physical ability to move.

PRESSURE ULCERS

MYTH OR STEREOTYPE

³ Tempkin, T., *Mobility: A Basic Human Need, Quality Care Advocate Special Section*, National Citizens’ Coalition for Nursey Home Reform, Washing DC, 1993, p.i.

⁴ Ibid.

⁵ Gueldner, S. H., and Spradler, J., “Outdoor Walking Lowers Fatigue,” *Journal of Gerontological Nursing*, Vol. 14, No. 10, pp. 6-12.

Because of the age-related changes in the skin and the frailty of nursing facility residents, pressure ulcers/sores are inevitable for individuals who are not independently mobile. Pressure sores are an unfortunate part of normal aging for frail, elderly persons.

REALITY⁶

A pressure ulcer is an injury caused primarily by unrelieved pressure that damages the skin and underlying tissue. An ulcer of this type is a serious problem that can lead to pain, longer hospital or nursing home stays, slower recovery from health problems, even death. Over 7% of residents in nursing facilities have pressure ulcers.⁷ Sixty percent (60%) or more of residents will typically be at risk of pressure ulcer development.⁸ Individuals who are at risk of developing pressure sores are those with limited mobility, incontinence, diabetes, decreased mental states, confusion, or apathy.⁹ *Almost all pressure ulcers can be prevented.*

The assessment of risk factors is critical to prevention and/or early detection and intervention. The primary risk factors are:

- immobility or unrelieved pressure, including pressure from use of a restraint,
- laying in urine or feces,
- poor nutrition and hydration.

All of the major causes can be addressed by facility staff and relate to basic, daily care routines.¹⁰

⁶ Most of this section is from: *Clinical Practice Guidelines No. 3: Pressure Ulcers in Adults: Predication and Prevention*, US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research, Rockville, MD, May 1992.

⁷ Harrington C., Carillo, H., Wellin, C., et al, *Nursing Facilities, Staffing, Residents and Facility Deficiencies*. University of California, San Francisco. August 2003.

⁸ Resident Assessment Protocol: Pressure Ulcers, Appendix C. HCFA's *RAI Version 2.0 Manual*. US Department of Health and Human Services, Health Care Financing Administration, Baltimore, MD. August 1999.

⁹ DiDomenico, D.L., and Ziegler, W.Z., *Positioning and Skin Care, Practical Rehabilitation Techniques for Geriatric Aides*, Aspen Publishers, 1989, p. 73.

¹⁰ *Ombudsman Guide to the Nursing Home Reform Law*, op.cit.

URINARY INCONTINENCE

MYTH OR STEREOTYPE

Urinary incontinence – the involuntary loss of urine – is to be expected, especially among residents in nursing facilities. It is another signal of advanced age and physical decline. Once it occurs, there is nothing that can be done except to keep individuals clean and dry.

REALITY

“Contrary to myth, incontinence is not a normal part of aging. It is actually easier to treat in the elderly than in the young. It is not inevitable, even in those with dementia (25% of the bedridden with dementia are continent), and is manageable in a third (33%) of those with dementia.”¹¹

Despite the high prevalence of urinary incontinence and the fact that it is associated with social and physical problems that impair general well being, nursing home staff often overlooks urinary incontinence as a potentially curable phenomenon.

It is estimated that more than one-half of all nursing home residents experience urinary incontinence. *Urinary incontinence is a symptom rather than a disease.*¹² In some cases, the disorder is temporary, secondary to an easily reversed cause such as a medication or an acute illness (e.g., urinary tract infection).¹³ The most probable cause of urinary incontinence is immobility caused by chemical or physical restraints or lack of a toileting program. In 2002 only 5.8% of residents in facilities had bladder training programs.¹⁴ Many cases are chronic, lasting indefinitely unless properly diagnosed and treated.¹⁵

“Despite the high prevalence of urinary incontinence and the fact that it is associated with social and physical problems that impair general well-being, nursing home staff often overlook urinary incontinence as a potentially curable phenomenon. Care plans that address incontinence often are custodial rather than rehabilitative in nature. In an attempt to keep residents dry, staff may diaper them, change clothing and linens frequently, toilet regularly, limit fluid intake, or use a catheter. Such approaches have their place under certain circumstances, but not until the resident has been evaluated properly to uncover the underlying cause of incontinence and treated when applicable.”¹⁶

“Continence depends on many factors. Urinary tract factors include a bladder that can store and expel urine and a urethra that can close and open appropriately. Other factors include the resident’s ability (with or without staff assistance) to reach the toilet on time (*locomotion*); his/her ability to adjust clothing so as to toilet (*dexterity*); cognitive function and social

¹¹ Siegal, D.L., “The Nursing Home Incontinence Project,” prepared for *Living is for the Elderly*, January 1992.

¹² Harrington, C., op.cit.

¹³ “Urinary Incontinence in Adults: Acute and Chronic Management.” Agency for Health Care Policy and Research, Public Health Service, Department of Health and Human Services Clinical Practice Guideline #2. 1996 Update.

¹⁴ Harrington, op.cit. 2003.

¹⁵ Burger, S. G., National Citizens’ Coalition for Nursing Home Reform, in a telephone conversation, January 14, 1994.

¹⁶ *Long Term Care Letter: Special Report: Incontinence*, Vol. 3, No. 9, Brown University, May 8, 1991.

awareness (*e.g., recognizing the need to void in time and in an appropriate place*); and the resident's motivation. Fluid balance and the integrity of the spinal cord and peripheral nerves will also have an effect on continence. Change in any one of these factors can result in incontinence, although alterations in several factors are common before incontinence develops.”¹⁷

In summary, incontinence not only affects skin conditions and care routines, but also has a profound effect on an individual's dignity, self-esteem, and social relationships. Minimizing risk factors and a thorough assessment and appropriate interventions are essential to helping individuals maintain, or regain, urinary continence. Restorative care is also important.

D E P R E S S I O N

MYTH OR STEREOTYPE

Older individuals tend to withdraw, slow down, and become depressed. Sadness is a natural response to loss of physical abilities and other life stage changes; therefore, depression is a normal part of living to an advanced age.

REALITY

“The ability to think, feel, interact with others, share a sense of purpose, work, love, experience gratification, care for others, and maintain self-responsibility are precious human attributes that elderly people strive to maintain. In only a few circumstances, are these elements of our experience and capacity so broadly and deeply challenged, as with depressive disease.”¹⁸

Depression in the elderly is being diagnosed and treated.¹⁹ A depressed mood may not be as noticeable a symptom among the elderly as are other symptoms such as loss of appetite, sleeplessness, lack of energy, and loss of interest and enjoyment of the normal pursuits of life. Depression affects many aspects of an individual's life. The risk of depression among women is over two times higher than that of elderly men.²⁰ One study suggests that a result of not treating depression in the elderly is a heightened risk of death.²¹ White men over 80 are at greatest risk for suicide of all older people.²² Treatment is effective, and depression can be alleviated in many cases. Proper assessment, detection, and intervention are critical.

¹⁷ Resident Assessment Protocol: Urinary Incontinence and Indwelling Catheter, Appendix C. HCFS's *RAI Version 2.0 Manual*. US Department of Health and Human Services, Health Care Financing Administration, Baltimore, MD August 1999.

¹⁸ *Diagnosis and Treatment of Depression in Late Life, Consensus Statement*, Vol. 9, No. 3, National Institutes of Health, Bethesda, MD, November 4-6, 1991.

¹⁹ Levenson, S., *Psychoactive Medications, Politics, The Unconventional "Wisdom" of LTC*, Caring for the Ages, February 2002.. In fact, Dr. Levenson says antidepressants are being overused without regard to the adverse effects that may accrue.

²⁰ Haight, B. and Hendrix, S., (1999) *Suicidal Intent/Life Satisfaction: Comparing Life Stories of Older Women*, *Suicide and Life Threatening Behavior*, 28(3) 272-284.

²¹ Golman, D., "High Death Risk is Found in Depressed Nursing Home Patients." *New York Times*, February 27, 1991.

²² Cronwell, Y., *Suicide in the Elderly*, in Schneider, LS., Reynolds, BD., Leowitz, BD., et al *Diagnosis and Treatment of Depression in Late Life: Results of a NIH Consensus Conference*, American Psychiatric Association Press 1994.

SAFETY CONCERNS

MYTH OR STEREOTYPE

As individuals become older and more physically frail, they need to be protected. Safety becomes very important; thus, minimizing risk is desirable. Using restraints is sometimes necessary to keep individuals from harming themselves by falling or other actions that may result in harm.

REALITY

All of life has risks. It is impossible to create a totally risk-free, 100% safe environment. However, some of the care practices that have been justified on the basis of safety may need to be questioned. “Physical restraints do not make people safer. In fact, restraints are often harmful. Caregiver experience and medical research now show:

Physical restraints do not make people safer.

When a person stops using a body part, that part no longer works very well. The old saying, ‘use it or you’ll lose it’ is true—people who are able to get up to try to walk and are restrained become weaker. Also, restrained residents often try to get out of restraints, sometimes resulting in serious injuries, such as broken bones, cuts requiring stitches, and concussions.

Some people also fall if they are not restrained. But research shows that these residents, when they do fall, have less serious injuries than those who are restrained.”²³

In talking with residents, families, and home staff, remember that individuals have the right to take risks and need enough information to allow them to make an informed decision. Advanced age does not remove an individual’s ability to accept risks. More information on restraints can be found in *Nursing Homes: Getting Good Care There*²⁴ or in the fact sheets on the web site of the National Citizens’ Coalition for Nursing Home Reform, www.nursinghomeaction.org.

²³ Burger, S.G. *Avoiding Physical Restraint Use: New Standards in Care: A Guide for Residents, Families, and Friends*. National Citizens’ Coalition for Nursing Home Reform. Washington, DC, 1993, p. 7.

²⁴ Burger, S.G., Fraser, V., Hunt, S., Frank, B., *Nursing Homes: Getting Good Care There*. Impact Publishers, Second edition, 2002. Available from the National Citizens’ Coalition for Nursing Home Reform, Washington, DC. (202)332-2275. www.nursinghomeaction.org.

APPENDIX D: Dementia

Dementia

“Dementia is a loss of mental function in two or more areas such as language, memory, visual and spatial abilities, or judgment severe enough to interfere with daily life. Dementia itself is not a disease but a broader set of symptoms that accompanies certain diseases or physical conditions.”¹

“The two most common forms of dementia in older people are Alzheimer’s disease and multi-infarct dementia (sometimes called vascular dementia). These types of dementia are irreversible, which means that they cannot be cured. In Alzheimer’s disease, nerve cell changes in certain parts of the brain result in the death of a large number of cells. Symptoms of Alzheimer’s disease begin to slowly and become steadily worse. As the disease progresses, symptoms range from mild forgetfulness to serious impairments in thinking, judgment and the ability to perform daily activities. Eventually, patients may need total care.

In multi-infarct dementia, a series of small strokes or changes in the brain’s blood supply may result in the death of brain tissue. The location in the brain where the small strokes occur determines the seriousness of the problem and the symptoms that arise. Symptoms that begin suddenly may be a sign of this kind of dementia. People with multi-infarct dementia are likely to show signs of improvement or remain stable for long periods of time, then quickly develop new symptoms if more strokes occur. In many people with multi-infarct dementia, high blood pressure is to blame. One of the most important reasons for controlling high blood pressure is to prevent strokes.”²

Before *dementia* became a common part of our vocabulary, the term *senility* was used. Senility and pre-senile dementia are still used as medical diagnoses. Regardless of the specific diagnosis, ombudsman approaches to residents with conditions that impair cognitive functioning are the same as described in this document.

*Conditions that can cause **reversible** dementia, if detected early, are:*


- Depression
- Drug interaction
- Problem with the thyroid gland
- High fever
- Minor head injury
- Poor nutrition
- Vitamin deficiency

¹ *Alzheimer’s Disease and Related Dementias Fact Sheet*. The Alzheimer’s Association. February 2004. (800)272-3900, www.alz.org

² *Forgetfulness: It’s Not Always What You Think*. Age Page. Alzheimer’s Disease Education and Referral Center. National Institute on Aging. US Department of Health and Human Services. <http://www.alzheimers.org/pubs/forgetfulness.html>.

Diseases that can cause irreversible dementia are:

- Alzheimer’s Disease
- Multi-Infarct Dementia or vascular disease caused by multiple strokes in the brain (MID)
- Parkinson’s Disease
- Creutzfeldt-Jakob Disease
- Huntington’s Disease
- Pick’s Disease
- Lewy Body Dementia



APPENDIX E: Alzheimer's Disease

*Alzheimer's Disease*¹²

Alzheimer's is:

- A disorder that destroys cells in the brain
- A degenerative, irreversible disease that usually begins gradually, causing a person to forget recent events or familiar tasks
- Variable in the rate with which it progresses from person to person
- Diagnosed as "probable Alzheimer's" based on a variety of tests. The diagnosis has an accuracy rate of 90%. Exact diagnosis can only be determined via a sample of brain tissues after death.

Residents have:

- Memory loss
- Confusion
- Personality and behavior changes
- Impaired judgment
- Difficulty communicating as the affected person struggles to find words, finish thoughts, or follow directions
- Inability to care for themselves as the disease progresses

Progression of Alzheimer's

Alzheimer's disease causes the formation of abnormal structures in the brain called plaques and tangles. As they accumulate in affected individuals, nerve cells connections are reduced. Areas of the brain that influence short-term memory tend to be affected first. Later, the disease works its way into sections that control other intellectual and physical functions.

Alzheimer's disease affects people in different ways, making it difficult for medical professionals to predict how an individual's disease will progress. Some experts classify the disease by stage (early, middle, and late). But specific behaviors and how long they last vary greatly, even within each stage of the disease.

As more is learned about the progression of the disease, new assessment scales are being developed to help physician's track, predict, and treat symptoms of Alzheimer's. New medications can slow the progression of memory loss in its early stages.

Statistics/Prevalence¹³

- Approximately 4.5 million Americans have Alzheimer's disease.
- 11 – 16 million Americans will have Alzheimer's by the middle of the next century unless a cure or prevention is found.

¹²*Facts: About Understanding Alzheimer's Disease.* The Alzheimer's Association, January 2004, op.cit.

¹³*Fact Sheet Alzheimer's Disease Statistics.* The Alzheimer's Association, April 2004. op. cit.

- One in 10 persons over 65 and nearly half of those over 85 have Alzheimer's disease. A small percentage of people in their 30s and 40s develop the disease.
- A person with Alzheimer's lives an average of 8 years but can live as many as 20 years or more from the onset of symptoms.

Understanding Behavioral Symptoms

Damage to the brain from Alzheimer's disease can cause a person to act in different or unpredictable ways. Some individuals with Alzheimer's become anxious or appear aggressive, while others repeat certain questions or gestures. Often these behaviors occur in combination, making it difficult to distinguish one from another. Behavioral symptoms do not always become apparent immediately after the onset of disease and often change as the disease progresses. Challenging behaviors not only cause discomfort to individuals with the disease, but also can be frustrating and stressful for caregivers who cannot understand them.

When behavioral symptoms surface, the individual first needs to be evaluated by a physician for potential treatable underlying causes. Behavioral symptoms often result from a variety of unmet needs or treatable problems that the individual cannot communicate, such as:

physical discomfort,

- medication side effects,
- chronic pain,
- infection,
- nutritional deficiencies,
- dehydration, or
- impaired vision or hearing.

When behavioral symptoms are brought on by causes other than physical problems, further evaluation should try to identify the unmet need and find ways to address it. Unmet needs include the basic human needs: need for toileting, sleeping, food, pain treatment, drink, warmth, companionship, and something useful to do. If a resident with dementia can no longer speak, behavior is the only form of communication.¹⁴

Non-Drug Treatments

Non-drug treatments of behavioral symptoms are recommended as a first option, since symptoms are best modified without the use of medication. Some suggestions for caregivers and families are:

- **Family education and counseling.** Learn what to expect when afflicted with or caring for someone with Alzheimer's. Family members who are familiar with the disease and know how to effectively communicate with their loved one may be able to better cope with behavioral symptoms. Counseling and support for individuals with the disease and their families is available through local chapters of the Alzheimer's Association.

¹⁴ *Ombudsman Guide to the Nursing Home Reform Law*. 2004. op.cit.

- **Modifying the environment.** Environmental factors such as lighting, color, and noise can greatly affect behavior. Dim lighting, for example, makes some individuals uneasy, while loud or erratic noise may cause confusion and frustration. The noise of a television set may be frightening. Modify the environment to reduce confusion, disorientation, and agitation. Keep familiar personal possessions visible to ensure comfort and feelings of warmth in your loved one's surroundings.

- **Planning activities.** The key to planning activities is in "knowing the details of a person's life."¹⁵ Help individuals with Alzheimer's organize their time and know what to expect each day. Planned activities help individuals feel independent and needed by focusing their attention on pleasurable or useful tasks. Daily routines such as bathing, dressing, cooking, cleaning, and laundry can be turned into productive activities and may be pleasurable for a housewife. Working on a motor for a mechanic, walking and gardening for a farmer are other examples. Other more creative leisure activities can include singing, playing a musical instrument, painting, walking, playing with a pet, or reading. Planned activities may relieve depression, agitation, and wandering, as well as help affected loved ones enjoy the best quality of life.

Drug Treatments

Non-drug treatments are not always effective; therefore, severe behavioral symptoms may be best treated with medication. In some cases, drugs that are available for the treatment of cognitive symptoms [such as donepezil HCl (Aricept®), or tacrine HCl (Cognex®)] also may improve behavioral symptoms.¹⁶

Several drugs are available for treating behavioral symptoms, and many more are being studied for specific use in helping individuals who suffer from Alzheimer's. Drugs commonly used to treat behavioral symptoms such as agitation, aggression, paranoia, delusions, or depression associated with Alzheimer's include:

Anti-psychotics (neuroleptics)

- Haloperidol (Haldol)
- Olanzapine (Zyprexa)
- Quetiapine (Seroquel)
- Risperidone (Risperdal)

Anti-anxiety drugs

- Alprazolam (Xanax)
- Buspirone (Buspar)
- Diazepam (Valium)

¹⁵ *Nursing Homes: Getting Good Care There.* 2002. *op.cit.*

¹⁶ *New drugs come on the market continually. While manufacturer's will claim they are safer than prior ones, that is rarely the case once the drug has been used in the general population for a time.* Sarah G. Burger, April 2004.

Antidepressants

- Amitriptyline (Elavil or Endep)
- Bupropion (Wellbutrin)
- Desipramine (Norpramin or Pertofrane)
- Fluoxetine (Prozac)
- Fluvoxamine (Luvox)
- Nortriptyline (Pamelor or Aventyl)
- Paroxetine (Paxil)
- Sertraline (Zoloft)
- Trazodone (Desyrel)

Like any other drugs, these treatments can cause undesirable side effects. Because individuals with Alzheimer's may have difficulty identifying medication side effects, caregivers should ask the physician or pharmacist about what to expect and warning signs to watch for with any drug that is prescribed. Key questions to ask about any medication are, "Does it *enable* an individual to function more independently or at a higher level? Does it *improve* an individual's quality of life?"¹⁷

Resources

The Alzheimer's Association is the only national voluntary health organization dedicated to research for the causes, cures, treatments and prevention of Alzheimer's disease and to providing education and support services to affected individuals and those who provide their care.

The Alzheimer's Association

225 N. Michigan Ave., Fl. 17 Chicago, IL 60601-7633
800-272-3900
www.alz.org

Greater Iowa Chapter

1730 28th Street
West Des Moines, IA 50266
515.440.2722 (P)
greateriowa@alz.org

East Central Iowa Chapter

317 7th Ave. SE, Ste. 402
Cedar Rapids, IA 52401
319.294.9699 (P)
319.294.0068 (F)

Midlands Chapter

705 North 16 Street
Council Bluffs, IA 51501
712.322.8840 (P)
402.502.7001 (F)

The Federal Government funds this service of the National Institute on Aging. It offers information and publications on diagnosis, treatment, patient care, caregiver needs, long term care, education and training, and research related to Alzheimer's disease. Staff responds to telephone and written requests and makes referrals to national- and State-level resources.

Alzheimer's Disease Education and Referral (ADEAR) Center

National Institute on Aging
Building 31, Room 5C27
31 Center Drive, MSC 2292
Bethesda, MD 20892
800-438-4380

¹⁷ Sarah G. Burger, consultant, April 2004.

APPENDIX F: Answer Key to Module Quiz

Correct answers are underlined

True or False

- T F 1. Depression in the older individuals is under diagnosed and under treated.
- T F 2. Medications prescribed for a medical problem may have an unintended effect on behavior.
- T F 3. It is possible for bones to break spontaneously.
- T F 4. Chronological age is an accurate indicator of an individual's feelings and abilities.
- T F 5. There are a range of non-drug interventions and treatments that may be effective in meeting a resident's needs when behavioral symptoms occur.
- T F 6. The possibility that a person will live in a care facility increases with age until age 78 when it remains the same no matter how long the person lives.
- T F 7. It is important to allow older people the extra seconds needed for their eyes to accommodate to changes in light or distance.
- T F 8. Individuals who have some degree of hearing loss may not realize that they have a loss.
- T F 9. One method of coping with change is through reminiscence.
- T F 10. Intelligence declines with the normal aging process.
- T F 11. "Once a man, twice a child," remains one of the great truths about the older.
- T F 12. Sensitivity to smells decreases with aging.
- T F 13. Sexual desire ceases in old age.
- T F 14. Movement is not important for individuals who are confined to bed.
- T F 15. Pressure sores are an unfortunate part of normal aging for frail, older persons.
- T F 16. Physical restraints prevent falls and injuries for individuals who are confused or have balance problems.
- T F 17. In spite of age related changes, individuals living in long-term care facilities are to be assisted in maintaining or improving their abilities unless a decline is unavoidable.
- T F 18. Inactivity increases calcium depletion which may contribute to osteoporosis.
- T F 19. Alzheimer's Disease affects areas of the brain that control long term memory first.
- T F 20. Incontinence is a normal part of aging.